

RAPID CREATION OF GENE TARGETING VECTORS USING HOMOLOGOUS RECOMBINATION IN YEAST

FIELD OF THE INVENTION

5 The present invention relates to the development of gene targeting vectors, i.e., vectors that are useful for causing directed mutation(s) in a target gene. More particularly, the preferred use of the gene targeting vectors of the present invention is for the mutation of genes in embryonic stem cells, wherein these mutated stem cells are then useful for the production of mutated whole animals, particularly mice.

BACKGROUND OF THE INVENTION

10 The Human Genome Project, which began in 1991, is expected to provide about 90% complete human genome sequence coverage by the year 2001. See e.g. Collins, *et al.*, Science, **282**:682-689, 1998. With this will be the discovery of tens of
15 thousands of new genes with unknown function. As detailed below, inactivation of genes in the mouse genome by homologous recombination is one of the best, albeit not one of the fastest, ways to assess function of a novel gene. In order to address this new demand for deciphering gene function it will be important to develop methods to make gene targeted mice faster and more efficiently. Using conventional methods
20 of making gene targeted mice, the majority of time is spent on building the targeting vector. Clearly, methods that allow a reduction in the time spent building gene targeting vectors will have a large impact on the total working hours invested in making gene targeted mice.

25 Mutations have long been a tool used to understand gene function. By observing the physical or biochemical defects that result when a gene is perturbed we can deduce, at least in part, the function of that gene. See e.g. Castle and Allen, Proc. Am. Acad. Arts Sci., **38**:603-621, 1903. Mutations can be found as spontaneous events in populations of organisms or induced with mutagens, such as the chemical mutagen N-ethyl-N-nitrosourea. Basic understanding of a mutation involves mapping
30 it or grouping it into a "complementation group" in order to define what gene a given mutation belongs to, or whether the mutation defines a novel gene. The process of mapping and genetic complementation testing involves breeding experiments, typically over several generations of the organism in question. For this reason, simple

organisms, such as the bacteria *Escherichia coli*, the yeast *Saccharomyces cerevisiae*, the round worm *Caenorhabditis elegans*, and the fruit fly *Drosophila melanogaster*, all with relatively short generation times, have been popular for experiments whereby novel mutations are induced through mutagenesis protocols and then characterized by genetic mapping and complementation testing. Until the late 1980's, mice, which are a popular organism for studying aspects of human physiology because of their relative physiological similarity, have not been as popular for studying genetics. Compared to the above-mentioned simple organisms, mice have a much longer generation time, and their size and husbandry requirements do not typically permit the rearing of the large numbers required in classical genetic experiments.

In late 1980's it became possible to create mutations in specific genes of interest in mice by the isolation of mouse embryonic stem cells (see e.g. Martin, Proc. Natl. Acad. Sci. USA, **78**:634-638; 1981, Evans and Kaufman, Nature, **309**:154-156, 1981), and the discovery that the phenomenon of "homologous recombination" could occur in mammalian cells (see e.g. Smithies, et al., Nature, **317**:230-234, 1985). These combined discoveries quickly led to a procedure that allowed one to inactivate a specific gene by first inactivating that gene in embryonic stem cells, and then transferring embryonic stem cells bearing the desired mutation into developing mouse embryos. See e.g. Doetschman, et al., Nature, **301**:576-578, 1987; Thomas et al., Cell, **219**:419-428, 1986. While this procedure is relatively laborious from the point of view of the work required to make a single mutation, it is efficient considering that a precisely defined mutation could be created in about 1 year using relatively few mice. Since these first demonstrations of targeted gene inactivation in mice, thousands of genes have been targeted and studied in this manner. The method has proven to be an invaluable tool for deciphering the complex functions of genes in the context of *in vivo* mouse biology. For example, the β -2-microglobulin gene was found to be necessary for the efficient development of the CD-4, CD-8⁺ class of T-cells, and consequently for the efficient production of certain classes of antibodies to certain antigens. See e.g. Koller, et al., Science, **248**:1227-1230; 1990; Spriggs, et al., Proc. Natl. Acad. Sci. USA, **89**:6070-6074, 1992. The connexin 43 gene, initially regarded as important in some of the very first cell differentiation events of mouse embryogenesis, was found to be most important in the developing heart. See e.g.

Reaume, *et al.*, *Science*, **216**:1831-1833, 1995. The JNK-3 gene was shown to be necessary in the cell death pathway in the brains of mice treated with the excitotoxin kanic acid. See. e.g. Yang, *et al.*, *Nature*, **389**:865-870, 1997.

Beyond just learning about the functions of genes, in some cases the mutant animals represent useful tools to study disease processes by allowing researchers to test potential therapeutic reagents. For example, U.S. Patent No. 5,777,195 describes a knockout of the mouse DARPP-32 gene and its utility as a screening model for potential therapeutic agents useful in the treatment of schizophrenia, Parkinson's disease, and the treatment of addictions, especially those involving drugs of abuse. U.S. Patent No. 5,814,300 describes gene-targeted heterozygous and homozygous SOD-1 null non-human mammals and their use for determining the effectiveness of a compound for counteracting the deleterious effects of oxidative stress. Another example may be found with U.S. Patent No. 5,859,307 which describes gene targeted mutants for the mouse RAG-1 gene and their potential use in the establishment of continuous lymphoid cell lines, as well as the establishment of animals developing tumors for use in studying tumor cell developments and treatments.

The method of making gene-targeted mutations in mice has proven particularly useful for the complex issue of deciphering the functions of individual family members in gene families. For example, there are 4 members of the prostaglandin E2 receptor family. Audoly, *et al.*, *Am. J. Physiol.*, **277**:H924-30, (1999) have shown that only the prostaglandin E2R EP1 member is responsible for perception of certain pain stimuli. Disruption of all 3 superoxide dismutase genes has shown that only Mn-superoxide dismutase is absolutely essential for survival. See e.g. Carlsson, *et al.*, *Proc. Natl. Acad. Sci.*, **295**:6264-6268, 1995; Reaume, *et al.*, *Natur. Genet.*, **13**:43-47, 1996; Li, *et al.*, *Nature Genet.*, **390**:376-381, 1995. Disruptions of all of the members of the src-kinase family have uncovered a diverse set of functions for each of the family members. See e.g. Lowell and Soriano, *Genes Dev.*, **10**:1845-1857, 1996.

The method of homologous recombination allows one to also make subtle changes in a gene without completely disrupting its function. The creation of a series of mutant versions of the gene that still possess partial activity has proven to be a complementary tool to completely inactivating a gene. See e.g. Nagy, *et al.*, *Curr.*

Biol.,8:661-664, 1998. In some instances mutations associated with human genetic disorders do not inactivate the gene but rather alter its function or constitute a "gain of function". Homologous recombination strategies that create more subtle changes can be employed in order to recapitulate these sorts of mutations in mice. For example, Sullivan, *et al.*, J. Biol. Chem.,272:17972-17980, (1997) have replaced the mouse apolipoprotein E gene with all 3 versions (alleles) of the human gene in order to more faithfully replicate the physiological effects associated with each of these 3 alleles. U.S. Pat No. 5,777,194 describes mice in which the A β portion of the mouse APP gene has been replaced with the human version of the gene in order to more faithfully replicate the physiology of A β deposition in the human brain, while providing a more sensitive assay with which to track A β levels.

SUMMARY OF THE INVENTION

This invention generally involves a process with which to create gene-targeting vectors in a rapid method by use of a specialized genomic DNA library, homologous recombination in yeast, and a subsequent selection system in *E. coli*. More particularly, this invention concerns a process with which to create gene-targeting vectors that may be used in mouse embryonic stem cells. The method involves using homologous recombination in yeast to simultaneously: 1) screen for genomic clones of interest and 2) build a gene targeting vector by virtue of replacing a defined portion of a gene of interest with a positive selection marker that can be used in both *E. coli* and mammalian cell culture.

In a first embodiment, the present invention provides a method of preparing a gene targeting vector, said method comprising transforming yeast cells with a RKO clone and a yeast targeting cassette (YTC), wherein said RKO clone comprises a genomic clone insert, a yeast replication element, a yeast selectable marker, a bacterial origin of replication, optionally a bacterial selectable marker, and optionally a mammalian negative selection marker, and wherein said YTC comprises a bacterial/mammalian positive selection marker flanked by recombinogenic arms, maintaining said yeast cells under conditions wherein said RKO clone and said YTC undergo homologous recombination via said genomic clone insert and said recombinogenic arms to produce a gene targeting vector, selecting transformed yeast cells by their expression of said yeast selectable

marker on said gene targeting vector or on said RKO clone, isolating said gene targeting vector and said RKO clone from said selected yeast cells, transforming bacterial cells with said gene targeting vector and said RKO clone, selecting transformed bacterial cells that grow on selective media that is selective for bacterial cells expressing said bacterial/mammalian positive selection marker, thereby selecting for bacterial cells transformed with said gene targeting vector, and isolating said gene targeting vector from said selected bacterial cells.

In a second embodiment, the present invention provides a method of preparing gene targeted mammalian cells having a targeted gene mutation, said method comprising transforming mammalian cells with said gene targeting vector of the first embodiment of the invention, maintaining said mammalian cells under conditions wherein said gene targeting vector and the genome of said mammalian cells undergo homologous recombination to produce a gene targeted mammalian cell, and selecting gene targeted mammalian cells wherein homologous recombination has occurred by selecting gene targeted mammalian cells for their expression of said bacterial/mammalian positive selection marker, thereby obtaining gene targeted mammalian cells containing said targeted gene mutation.

In a third embodiment, the present invention provides a method of making gene targeted mice, said method comprising combining a gene targeted mouse cell according to the second embodiment of the invention with an early mouse embryo to produce a gene targeted embryonic construct, and introducing said gene targeted embryonic construct into a female host mouse, wherein said gene targeted embryonic construct is allowed to mature into a chimeric live whole mouse, said whole mouse thereby having a genome that includes said targeted gene mutation.

In a fourth embodiment, the present invention provides a gene targeting vector comprising a yeast replication element, a yeast selectable marker, a bacterial origin of replication, optionally a bacterial selectable marker, optionally a mammalian negative selection marker, and a genomic clone insert containing a bacterial mammalian positive selection marker inserted therein.

DESCRIPTION OF THE FIGURE SHEETS

FIGURE 1 provides a general paradigm for introducing mutations into a mammalian genome using homologous recombination. In the figure, homologous

recombination between elements A and B results in a product C wherein region 3 has been replaced by a marker segment (Neo). The negative selection marker (Hsv-TK) is separated from the positive selection marker in the process of homologous recombination.

5 FIGURE 2 provides a plasmid map of a preferred RKO backbone according to the present invention. ARS1 is a yeast autonomous replicating sequence that serves as a yeast replication element. CEN4 is a yeast centromeric sequence that helps to stabilize propagation of the vector in yeast. TRP1 is a yeast selectable marker. Ori is a bacterial origin of replication. Amp is a bacterial selectable marker.
10 The hsv-TK cassette is a mammalian negative selection marker. Addition of the genomic clone insert to this plasmid would yield an RKO clone. In a preferred embodiment, genomic clone inserts would be inserted into the multiple cloning site (MCS) shown in this plasmid to create the RKO clone. Addition of Cos sites allow the RKO clone to be packaged in lambda phage as a cosmid clone for propagation
15 purposes.

 FIGURE 3 depicts an example of a preferred yeast-targeting cassette (YTC) as described in this invention. The bacterial/mammalian selection marker cassette is flanked by recombinogenic arms (RA). A promoter (T7) allows transcription of the bacterial/mammalian positive selection marker (Neo^r) to occur in bacteria. An
20 internal ribosome entry site (IRES) element allows translational initiation of the said selection marker to occur when the said marker is transcribed by a promoter in the genomic region of interest (GRI). A Shine-Dalgarno element (SD) allows translation of the said selection marker to occur in bacteria. Fusion of the coding sequence for the said selection marker with the coding sequence for a reporter protein such as
25 bacterial beta galactosidase (lacZ) creates a reporter system for the GRI so that it may be easily possible to determine when a gene of interest is being expressed. A polyadenylation signal (pA) is desirable if transcription in mammalian cells initiates from the (GRI).

30 **DETAILED DESCRIPTION OF THE INVENTION**

Definitions

Arms of Homology: A pair of DNA segments that is present in the gene targeting vector; these segments are homologous to two portions of the target gene. Because of this homology, the arms of homology will be able to undergo homologous recombination with the target gene. The arms of homology are preferably at least 200
5 bp in length to allow for appreciable rates of homologous recombination to occur in mammalian cells. More preferably, the arms of homology will be at least 2 kb or greater in length.

Bacterial Origin of Replication: A genetic element that enables replication of the DNA sequence of which it is a part during each cell cycle in a bacterium, such
10 that the DNA sequence will be maintained in each daughter cell. Examples include pMB1, pSC101 and pUC.

Bacterial Selectable Marker: A genetic element encoding a protein that when expressed in bacteria enables selection of the bacteria by the presence of the protein. Thus, any bacteria that contain and express a bacterial selectable marker
15 can be differentiated from otherwise similar bacteria that do not contain and express the marker. Examples include Amp^r, Tet^r, and Kan^r.

Bacterial/Mammalian Positive Selection Marker: A genetic element encoding a protein that when expressed in a bacterial or mammalian cell enables selection of the cell for the presence of the protein. In an appropriate selective
20 environment, only cells expressing the marker protein should survive and multiply. Thus, any cell that contains and expresses a bacterial/mammalian positive selection marker can be differentiated from otherwise similar cells that do not contain and express the marker. Examples include Neo^r, Zeo^r and *bsd*.

Early Mouse Embryo: A mouse embryo that is capable of being combined
25 with a gene targeted mouse cell as defined herein and developing into a chimeric adult mouse. Preferably, an early mouse embryo has developed 2.5 days *post-coitus* (dpc) to 3.5 dpc, which is approximately equivalent to the 8-cell stage through the blastocyst stage in development.

Embryonic Stem Cells: Stem cells of embryonic origin. Preferably,
30 embryonic stem cells are derived from embryos that have developed to 3.5 dpc, which is approximately equivalent to the blastocyst stage in development.

Gene Targeted Cell: A cell containing a non-naturally occurring genomic mutation as a result of homologous recombination between the cell genome and a gene targeting vector as defined herein.

Gene Targeted Embryonic Construct An embryo resulting from the combination of an early mouse embryo with a gene targeted cell. When introduced into a female host mouse, a gene targeted embryonic construct is capable of developing into a chimeric live whole mouse.

Gene Targeting Vector: A DNA molecule that results from homologous recombination between a RKO clone and a YTC as defined herein. A gene-targeting vector contains a mutated target gene that is ready to knockout or otherwise mutate its counterpart gene via homologous recombination when inserted into a cell containing that counterpart gene.

Genomic Clone Insert A DNA segment that originates from the genome of a particular organism of interest. A genomic clone insert is part of a RKO clone as defined herein. Typically, this segment is derived from the creation of a library of clones from a genome. One aspect of the present invention is the selection of a specific desired genomic clone insert during the process of creating the gene-targeting vector. However, at the beginning of the invention method, RKO clones are produced that contain numerous genomic clone inserts, not only the desired one, and thus this term encompasses all such DNA segments derived from the genomic library.

Genomic Region of Interest (GRI) A contiguous sequence of genomic DNA comprising greater than 100 base pairs and which one wants to specifically alter using the present invention. Typically the GRI will include the coding portion of a gene of interest but could also include a regulatory region that one wants to alter in a specific way. Still another example involves an intervening sequence between a gene(s) of interest that one may want to alter in some way in a multistep process, part of which involves creating specific alterations using the present invention.

Homologous Sequence: A sequence that is at least about 90% identical to the comparison sequence, and is preferably greater than 95% identical. Sequence identity, as used throughout the present application and claims, is defined using the BLAST software for making sequence comparisons.

Mammalian Negative Selection Marker. A genetic element that when present in a mammalian cell enables selection of the mammalian cell for the absence

of the element. In an appropriate selective environment, any cells expressing the marker protein will not survive and/or multiply. Thus, any mammalian cell that contains a mammalian negative selection marker can be differentiated from otherwise similar cells that do not contain and express the marker. Examples include Herpes Simplex Virus-Thymidine Kinase (HSV-TK), Diphtheria toxin (DT), and a promoter that makes antisense RNA designed to inactivate a positive selection marker (Skryabin and Schmauss, Transgenic Res., 6:27-35, 1997). Examples of negative selection markers useful in the present invention are described in U.S. Patent No. 5,464,764.

Multiple Cloning Site: A DNA segment that is optionally present in the RKO clone; this segment comprises a series of very closely situated restriction enzyme recognition and cutting sites. In fact, the sites are frequently adjacent and possibly even overlapping.

Mutation: A change in the sequence of a DNA segment, such that the sequence is no longer identical to a naturally occurring sequence.

Recombinogenic Arms (RAs): A pair of DNA segments that are present in the YTC; these segments are homologous to two portions of a target gene. Because of this homology, the recombinogenic arms will be able to undergo homologous recombination with the target gene. The RAs are preferably at least 20 bp in length to allow for appreciable rates of homologous recombination to occur in yeast. More preferably, the RAs will be at least 100 bp in length.

RKO Backbone: This is the RKO clone without the genomic clone insert. Because this vector is grown in bacterial cells to create and manipulate it, a bacterial selectable marker is typically present, and therefore still present in the RKO clone. However, the presence of this marker is unnecessary for practicing the methods of the present invention.

RKO Clone: A DNA molecule comprising a genomic clone insert, a yeast replication element, a yeast selectable marker, a bacterial origin of replication, optionally a bacterial selectable marker, and optionally a mammalian negative selection marker, each as defined herein. The RKO clone may include additional genetic elements as well. RKO is an acronym for RapidKnockOut, which is the terminology used to describe the preferred use of the present invention, for rapidly building a gene-targeting vector. Homologous recombination in yeast between a YTC and an RKO clone yields a gene-targeting vector.

Stem Cells: Pluripotent or totipotent cells that are capable of being combined with an early embryo and subsequently maturing into a chimeric adult.

Substantially Isogenic: A sequence that is at least about 97-98% identical to the comparison sequence, and preferably is at least about 99% identical.

5 **Targeted Gene Mutation:** A mutation created in the GRI by practice of the present invention. The mutation is created by virtue of homologous recombination occurring between the RAs(recombinogenic arms) of the YTC and the genomic clone insert of the RKO clone so as to create a gene-targeting vector. The targeted gene mutation is subsequently transferred into the genome of the target cell and/or
10 organism by homologous recombination between the gene-targeting vector and the endogenous GRI within said target cell.

15 **Yeast Replication Element** A genetic element that enables replication of the DNA sequence of which it is a part during each cell cycle in yeast, such that the DNA sequence will be maintained in each daughter cell. Examples include 2 μ origin of replication, a yeast autonomous replicating sequence (ARS) including those
sequences which may be found in the genomes of other organisms that allow replication in yeast (Larionov, *et al.*, Proc. Natl. Acad. Sci. USA, **93**:491-496, 1996) .

20 **Yeast Selectable Marker.** A genetic element encoding a protein that when expressed in yeast that enables selection of the yeast cell by the presence of the protein. Thus, any yeast cell that contains and expresses a yeast selectable marker can be differentiated from otherwise similar yeast cells that do not contain and express the marker. Examples include TRP1, HIS3, URA3, and LEU2.

25 **Yeast Targeting Construct (YTC):** A DNA molecule comprising a bacterial/mammalian positive selection marker flanked by recombinogenic arms, each as defined herein. The YTC may comprise additional genetic elements as well. In a preferred embodiment of this invention the bacterial/mammalian positive selection marker is constructed in such a way that, while capable of being expressed in bacteria under all normal growth conditions, it is only expressed in mammalian cells if it is
30 inserted within a gene of the mammalian cell. For example, if the bacterial/mammalian positive selection marker has no mammalian promoter but has a bacterial promoter and an internal ribosomal entry site (IRES) element, then mammalian positive selection will only occur if the bacterial/mammalian positive

selection marker is inserted within a gene that is being expressed. Alternatively, if the bacterial/mammalian positive selection marker lacks a polyadenylation signal then again mammalian positive selection will only occur if the marker is inserted within a gene that has a polyadenylation signal (as should be the case in most gene-targeting experiments).

Figure 1 depicts the general paradigm for introducing mutations into a mammalian genome using homologous recombination (reviewed in Capecchi, Trends Genet, **217**:70-76, 1989; and Koller and Smithies, Ann. Rev. Immunol, **218**:705-30, 1992). This general paradigm is applicable to the methods of the present invention, as well as to prior art methods. A length of genomic DNA is first depicted by organizing it into regions (numbered 0-6 in FIG. 1, element A). In FIG. 1, region 3 is designated to be deleted. Homologous recombination using a gene-targeting vector is utilized. The type of gene-targeting vector used for the deletion of a gene is termed a replacement or deletion vector. "Deletion vector" as used herein refers to a vector that includes one or more selectable marker sequences and two sequences of DNA homologous to the genomic DNA that flank the DNA gene sequence to be deleted. These flanking sequences are termed arms of homology. In FIG. 1, element B, the arms of homology are represented by regions 1-2 and 4-5. Preferably, these arms of homology are substantially isogenic for the corresponding flanking sequences in the cell being targeted or "target cell." The use of DNA that is substantially isogenic to the target cells DNA helps assure high efficiency of recombination with the target sequences (See U.S. Pat. No. 5,789,215). The deletion vector preferably includes at least a positive selection marker within the arms of homology to enable the scoring of recombination. Such positive selection markers can confer a phenotype not normally exhibited by wild-type mammals; for example, resistance to a substance normally toxic to the target cells. In another example, the deletion vector also includes one or more negative selection markers outside the arms of homology to facilitate identification of proper homologous recombinants. U.S. Patent No. 5,464,764 describes the use of such "positive-negative" selection methods. The negative selection markers can confer, for example, sensitivity to a substance not normally toxic to the target cell. The selection markers can be gene cassettes, i.e., include both a promoter and an accompanying coding sequence. The result of homologous

recombination of the gene-targeting vector with cellular DNA in the paradigm is shown in FIG. 1, product C. As depicted therein, region 3 has been replaced by the positive selection marker. Upon successful gene-targeting and homologous recombination, the positive selection marker is incorporated into the genome within the arms of
5 homology in place of the deleted gene segment, while the negative selection marker is excluded. Thus, to enrich for homologous recombinants, gene-targeted cells are grown in culture medium containing the appropriate positive and negative selective compounds.

The "target cells" are those cells to be mutagenized. The target cells preferred
10 for use in the present invention are mouse embryonic stem cells. However, other cells can be utilized. If embryonic stem cells are to be targeted then one can utilize properly identified gene targeted cells to combine with early mouse embryos to create a gene targeted embryonic construct. Preferred methods for combining gene-
15 targeted cells with early embryos include blastocyst injection and aggregation methods. The gene targeted embryonic construct is then transferred to a host animal, preferably a pseudopregnant female mouse. Some of the resulting offspring mice will mature and produce gametes that bear the desired mutation. These particular mice are referred to as "germline chimeras". Some of the offspring of germline chimeras bear one mutant copy of the gene of interest in all of the cells in their body and are
20 referred to as "heterozygotes". If two of these heterozygotes are mated to each other some of their offspring will bear two mutant copies of the gene of interest and are referred to as homozygotes.

The method described herein uses a genomic DNA library made in a vector with the following features:

- 25 a yeast replication element;
- a yeast selectable marker;
- a bacterial origin of replication;
- optionally a bacterial selectable marker; and
- optionally a negative selection marker
- 30 to which is added a genomic clone insert. An example of such a vector is shown in figure 2.

If the library is to be a cosmid library then at least one Cos site must be included. In a preferred embodiment, the vector includes two Cos sites in the same

orientation in tandem repeat with a unique restriction site separating the two. For a thorough description on the use of Cos sites to build cosmid libraries see Maniatis, *et al.*, Cold Spring Harbor, New York, (1982), incorporated herein by reference.

In another preferred embodiment, a region adjacent to the negative selection
5 marker is a multiple cloning site containing a large number of rare cutters (e.g., 6 or more base pairs in the recognition sites). Preferably these sites are strategically placed. For example a *Bam*HI site, which is often used as a recipient site to receive genomic DNA that has been partially digested with the enzyme *Sau*3A (see Maniatis, *et al.*, *supra*), would preferably be placed in a center position relative to
10 the total collection of enzyme sites in this region.

A genomic DNA library may be constructed making a RKO clone with the features described above. For the purposes of making gene-targeting vectors in mammalian cells, the source of DNA is preferably from an isogenic source. Such
15 would be the case if mouse cells were to be targeted and genomic DNA used to make the gene targeting vector was isolated from the same strain of mice. Further description on the use of isogenic DNA can be found in U.S. Pat. No. 5,789,215. Methods used to make genomic DNA libraries are described in Maniatis *et al.*, *supra*.

A preferred embodiment of this invention involves screening a genomic
20 library comprising the above features by combining a DNA library (an aliquot of DNA representing either the entire library of recombinant genomic clones or a subset of the library) with a fragment of DNA called the yeast targeting cassette (YTC). A DNA aliquot representing the entire library can be obtained by inoculating a culture with a number of colony forming units or plaque forming units equal to or
25 greater than the total number of independent genomic clones known to be in the library. An aliquot of DNA appropriately prepared from this culture would then represent the entire genomic library. Similarly if the culture was initially inoculated with fewer clones than what represented the entire library then DNA prepared from this culture would represent a subset of the entire library.

30 A preferred method of screening the library involves dividing the library into pools of recombinant clones representing subsets of the library. All of the pools are screened by either conventional polymerase chain reaction (PCR) methods designed to identify a GRI or by hybridizing radioactively labeled probes to samples

of the pools that have been blotted onto a suitable membrane. Clones bearing the GRI are then further isolated by mixing them with the YTC and performing the procedures described herein.

An example of a preferred YTC is shown in figure 3. The YTC comprises a
5 bacterial/mammalian positive selection marker flanked by regions of sequence identity to the GRI. These regions of sequence identity will serve as recombinogenic arms. The bacterial/mammalian positive selection marker can be flanked by loxP or FRT sites, thereby allowing for excision of the bacterial/mammalian positive selection marker (see, for example, U.S. Pat. No.
10 4,959,317). In a preferred embodiment of this invention the bacterial/mammalian positive selection marker is constructed in such a way that, while capable of being expressed in bacteria under all normal growth conditions, it is only expressed in mammalian cells if it is inserted within a gene of the mammalian cell. For example, if the bacterial/mammalian positive selection marker has no mammalian promoter but
15 has a bacterial promoter and an internal ribosomal entry site (IRES) element, then mammalian positive selection will only occur if the bacterial/mammalian positive selection marker is inserted within a gene that is being expressed. Alternatively, if the bacterial/mammalian positive selection marker lacks a polyadenylation signal then again mammalian positive selection will only occur if the marker is inserted within a
20 gene that has a polyadenylation signal (as should be the case in most gene targeting experiments).

In a further preferred embodiment of this invention the YTC will bear a reporter system capable of being expressed when inserted within the GRI. For example, if the bacterial/mammalian positive selection marker is to be expressed in
25 mammalian cells by the GRI by virtue of an IRES element then the bacterial/mammalian positive selection marker may be a fusion protein between a reporter protein (such as betagalactosidase) and a selection protein (such as neo'). Alternatively, if the bacterial/mammalian positive selection marker is not expressed by the GRI, then the reporter protein can be expressed by the GRI by virtue of an
30 IRES element that allows reporter protein translation to occur from mRNA transcripts driven by the native promoter in the GRI.

A preferred embodiment of this invention involves creating the YTC by use of a single PCR reaction using 2 chimeric oligonucleotides. The 3' portion of each

oligonucleotide bears sequence identity to the selected mammalian positive selection marker. The 5' portion of each oligonucleotide bears sequence identity to the GRI. A conventional PCR reaction is performed using these two oligonucleotides and a plasmid bearing the bacterial/mammalian positive selection marker as a template. The resulting amplified product will be a bacterial/mammalian positive selection marker flanked by RAs, thereby forming the YTC.

The RAs are meant to allow homologous recombination to occur in yeast between genomic clones bearing the GRI (i.e., a RKO clone) and the YTC. Figure 2 illustrates aspects of the YTC and its regions of sequence identity with the GRI. Both RAs should be derived from the same chromosomal region of the GRI and must be oriented in the same direction and order as they are positioned in their endogenous chromosomal location. See e.g. WO 93/03183. They may either represent contiguous elements in the genome from which they are derived or be spaced by a gap. Preferably any gap is less than 2 million base pairs in size.

Following the step in which the YTC is mixed with library DNA, the mixture is transfected into a yeast strain bearing a mutation in a gene that is complemented by the yeast selectable marker. The transfected yeast are grown on minimal medium agar plates (selective conditions). All of the colonies that grow on these plates represent individually transfected yeast cells. Although not selected-for at this stage, yeast colonies that bear a genomic clone of interest and were also transfected by the YTC are susceptible to undergoing homologous recombination between the GRI and the YTC by virtue of the RAs flanking the YTC. Figure 1 illustrates the process of homologous recombination between the YTC and the GRI. Note that only yeast cells receiving a genomic clone bearing the GRI will undergo homologous recombination with the YTC and thereby incorporate the positive selection marker associated with the YTC. Also note that if the RAs on the YTC represent contiguous elements from the gene of interest then no deletion within the GRI occurs. However, if the regions of sequence identity represent noncontiguous elements from the GRI then the result of homologous recombination in yeast is a deletion of the entire region between the corresponding RAs. This latter strategy is useful for inactivating a gene of interest.

Depending on the complexity of the library aliquot that was used in the previous step, an appropriate number of yeast colonies are scraped off the agar plates. For example, if the library aliquot used represented 1,000 different genomic clones then 500 times this number, or 500,000 yeast clones, would preferably be scraped off of the agar plates. If the library aliquot represented 100 genomic clones, then 500 times this number, or 50,000 yeast clones, would preferably be scraped off of the agar plates.

The yeast clones that were scraped off would then preferably be subjected to one of several methods available to shuttle plasmids from yeast to bacteria such as the "smash and grab" procedure or Zymoprep™ (Zymo Research, Orange, CA) method. Briefly, the former method involves sonicating or vortexing the yeast cells in the presence of glass beads and phenol in order to crudely extract plasmid DNA. The crude DNA extract is then transfected into *E. coli* by conventional bacterial transformation protocols.

The transfected *E. coli* are then plated onto nutrient rich agar plates containing a drug that selects for the positive selection marker present in the YTC. Only genomic clones that bear the GRI and underwent homologous recombination with the YTC in yeast should bear a stable copy of the marker present in the YTC.

In this way genomic clones bearing the region of interest are isolated and simultaneously manipulated in order to create a gene-targeting vector for mammalian cells. Note that the product that is created by homologous recombination in yeast bears two arms of homology in the correct orientation, separated by a positive selection marker and flanked on one side by a negative selection marker. Note that the relatively small recombinogenic arms of the YTC have essentially been increased in size by merging with the larger genomic clone insert of the RKO clone, thereby creating arms of homology, which are typically much larger than recombinogenic arms. RAs can be relatively small, as shorter segments will recombine in yeast at acceptable rates, whereas relatively longer homologous sequences are required for homologous recombination to occur at acceptable rates in mammalian cells.

Once the gene-targeting vector has been created, it is then preferably used to prepare gene targeted mammalian cells. Preferably, these cells are mouse cells, and can later be used to create a mouse carrying a targeted gene mutation.

First, mammalian cells are transformed with the gene-targeting vector by any method. There are numerous methods known in the art, and many more will likely be developed. Which method is used is not deemed critical to the practice of the present invention. After transformation, the cells are maintained under
5 conditions that will allow gene-targeting vector to undergo homologous recombination with the GRI, thereby producing a gene targeted mammalian cell. Cells wherein homologous recombination has occurred may be selected by their expression of the bacterial/mammalian positive selection marker, which will have been inserted into the genome by the recombination event. Additionally, if the gene
10 targeting vector included a mammalian negative selection marker outside the arms of homology, then the transformed mammalian cells wherein homologous recombination have occurred may be selected by their non-expression of this negative marker.

It is preferred that the mammalian cells used be stem cells. Even more
15 preferred is the use of embryonic stem cells. However, any cell in which it would be desirable to create a targeted gene mutation may be used.

To make a gene targeted mouse, the gene-targeted cell as described above is used to grow a new whole live mouse. Preferably, this is done by taking the gene targeted mouse stem cell and combining it with an early mouse embryo to produce
20 a gene targeted embryonic construct. What you will then have is an early mouse embryo that includes at least one cell carrying the targeted gene mutation as desired. Next, the gene targeted embryonic construct is introduced into a female host mouse, preferably a pseudo-pregnant female host mouse. The gene targeted embryonic construct will then be able to implant itself in the host mouse uterus, and
25 mature in normal fashion into a live baby mouse. This baby mouse will be chimeric, with some cells derived from the early mouse embryo cells, and others derived from the gene targeted mouse stem cell. When this mouse is bred to other regular mice, some of their offspring will be heterozygotes whose genome includes the gene-targeted mutation.

30 It is frequently desirable to produce mice that are homozygous for the gene-targeted mutation. This is done by merely crossbreeding heterozygotes until a homozygote offspring is born. In some cases, the mutation will be lethal, and it will

not be possible to obtain live homozygous mice. In other cases, it will be possible but extremely difficult.

The present invention is described in more detail with reference to the following non-limiting examples.

5

EXAMPLES

EXAMPLE 1 - Building of pCosRKO Cosmid Library Backbone Vector

The CEN4, ARS1, TRP1, Amp^r, and pUC ori elements of YCplac22 (Gietz and Sugino, Gene, **74**:527-534, 1988; obtained from American Type Culture Collection, 10801 University Boulevard, Manassas, VA) were amplified by polymerase chain reaction (PCR) (Mullis and F.A., Methods Enzymol., **155**:335-350, 1987) using primers 44413-12A (CGC TTC GAA ATC GCT GGG CCA TTC TCA TGA; SEQ ID NO:1) and 44413-12B (CGC ATT TAA ATT CTT CCG CTT CCT CGC TCA CT; SEQ ID NO:2) which hybridize to the 5' end of the CEN4 domain and the 3' end of the pUC ori respectively. This PCR product was digested with the restriction endonucleases *Bst*BI and *Swa* I to create sticky ends. Similarly the 2 Cos sites of plasmid SuperCos (Stratagene Inc., La Jolla, Calif.) were amplified using primers 44413-12C (GAC CGC TTA TCA TCG ATA AGC; SEQ ID NO:3) and 44413-12D (CGC ATT TAA ATG AAG GCT CTC AAG GGC ATC GG; SEQ ID NO:4). This PCR product was digested with the restriction endonucleases *Swa*I and *Cla*I to create sticky ends. The two digested PCR products were purified using agarose gel electrophoresis and ligated using T4 ligase. The ligation mixture was used to transform competent *E. coli* cells, and clones were identified that represented the correct product carrying the appropriate fragment of YCplac22 fused to the 2 Cos sites from pSuperCos. This construct was called YCpCos.

YCpCos was PCR amplified with primers 44413-12E (CGC GGA TCC TAG CCT GGC CAT CCG GGC CAT CGC TGG GCC ATT CTC ATG A; SEQ ID NO:5) and 44413-12F (CGC ATC GAT TTC GAT AAG CTC TGC TTT TTG T; SEQ ID NO:6). This PCR product was digested with the restriction endonucleases *Bam*HI and *Cla* I to create sticky ends. Similarly, the herpes simplex virus thymidine kinase (hsvTK) expression cassette was amplified from the plasmid pJNS using the primers 44413-12G (CGC TTC GAA CTA CCG GGT AGG GGA GGC GCT; SEQ ID NO:7) and 44413H (CGC AGA TCT TCC GAT GGC CGC AGC GGC CTC TGA

TGG AAT TAG AAC TTG GCA; SEQ ID NO:8). This PCR product was digested with the restriction endonucleases *Bst*BI and *Bgl*II to create sticky ends. The two digested PCR products were purified using agarose gel electrophoresis and ligated using T4 ligase. The ligation mixture was used to transform competent *E. coli* cells, and clones were identified that represented the correct product carrying the hsvTK cassette fused into YCpCos. The construct was called YCpCosTK.

Finally, the multiple cloning site (MCS;

GGCCGCTGCTTCCAGGCTAACACTTATATAGGCATTGGTACTGAGATACAGCG
AGATACCGTGTAACCTATAACGGTCCTAAGGTAGCGAAGGCTAACACTTATATAG
GCATTGGTACTGAGATACAGCGAGATACCGTGCGGCCGCTGCGGCCAATTAA
CCCTCACTAAAGGGAGGCGCTATGCATACTAGTCCATGGGATATCCCCGGGGTT
AACCCGCGGCTCGAGATCGATGAATTCGGCCGGCCGTTTAAACGGATCCGGC
GCGCCTTAATTAAAAGCTTCATATGTCTAGAGTCGACGAGCTCGGTACCGGGC
CCGCATGCTGGCCAAGGCCTAGATCTCCCTATAGTGAGTCGTATTATGGCAAA
CAGCTATTATGGGTATTATGGGTAGGCCATCCGGGCGCGGCCGCTTCCATC
CGGGCC; SEQ ID NO:9) was created using a series of overlapping and
complementing oligonucleotides and cloned into YCpCos by linearizing YCpCos
using the restriction endonuclease *Sfi*I. The resulting plasmid was called
pCosRKO.

EXAMPLE 2 - Building a Genomic Library with pCosRKO

Mouse genomic DNA was isolated from one kidney of a female DBA/1lacJ mouse using a standard proteinase K digestion followed by phenol/chloroform extraction and ethanol spooling of the DNA. Briefly, the kidney was bisected approximately 10-15 times with sterile scissors. The minced kidney was incubated in 5 ml of a tissue digestion buffer (50mM Tris Cl (pH 8.0) , 100 mM EDTA (pH 8.0), 100mM NaCl, 1% SDS, 600 µg/ml proteinase K (Roche, Cat. No. 1 373 196)) overnight at 55°C. The digestion was then extracted with an equal volume of a 1:1 phenol/chloroform mixture. DNA was then precipitated by adding two volumes of ethanol and spooling the DNA using a sterile glass rod. Finally, spooled DNA was scraped off of the glass rod into a 1.5 ml tube and resuspended in 300 µl of TE (10 mM Tris Cl (pH 7.5) 1mM EDTA) . Quality of the isolated DNA was checked by

running uncut DNA on a pulse field agarose gel apparatus along with molecular weight markers (Low Range PFG Marker; New England Biolabs® Inc., Beverly, MA). Specifically, a CHEF-DR® II apparatus (BioRAD, Hercules, CA) was used with a 1% agarose gel, 1-6 second switch time, ramped over 11 hours, and 6 V/cm at 120° angle. The gel was run for 11 hours. Good quality DNA was at least 150 kb or larger. A series of partial digests of the genomic DNA with the enzyme *Sau3A* were performed by mixing 15 µl of DNA with 1.5 units of enzyme (Boehringer Mannheim) in a 100 µl reaction volume. The mixture was preheated to 37°C before adding enzyme. 15 µl aliquots were removed at 0, 1, 2, 3, 4, 8, and 15 minutes after adding enzyme. Immediately after removing an aliquot 3 µl of 0.5M EDTA was added to the aliquot and the mixture was placed at 65°C for 10 minutes. The aliquots were loaded onto a pulse field gel apparatus and run as described above. It was determined that a 2.5 minute digest time was ideal as it generated a mean fragment length that was closest to the preferred 30 – 40 kb length. A large scale *Sau3A* digest was performed by mixing 150 µl of DNA with 15 units of enzyme in a 1000 µl reaction volume. Again, the mixture was preheated before the enzyme was added. The 1 ml reaction was loaded onto a 17 ml 40%-10% sucrose gradient (see Maniatis *et al.*, *infra*) and spun at 26K rpm for 24 hours at 15°C. Every third aliquot was ethanol precipitated, and resuspended in 50 µl of TE (10 mM Tris (pH 7.5), 1 mM EDTA). 10 µl of each TE-resuspended aliquot was run on a pulse field gel with markers as described above. From this analysis it was inferred which aliquots contained mostly fragments that were between 30 and 40 kb. The 30-40 kb-prevalent aliquots were pooled.

PCosRKO vector DNA was prepared by digestion with *PvuII*, treatment with calf intestinal phosphatase, and then digestion with *Bam*HI. Individual vector arms were isolated on a 0.7% agarose TBE gel using Qiagen® gel extraction kit (Qiagen, Valencia, CA). One µg of the double digested pCosRKO vector was ligated to 4.0 µg of the size selected *Sau3A* genomic DNA in a 10 µl reaction volume. Ligation was performed at 4°C overnight. Then, the ligation reaction 1 µl of the ligation mixture was packaged using Gigapack Gold packaging extract (Stratagene, LaJolla, CA).

Following the 2 hour packaging reaction, the titer of the library was determined by diluting the packaging reaction 1:10, 1:50, and 1:100 in SM buffer, adsorbing phage to properly prepared XL-1 Blue MRA cells (Stratagene, La Jolla, CA), and plating on LB-ampicillin plates as described by the packaging extract manufacturer (Stratagene, La Jolla, CA). Numbers of colonies from each of the dilutions was determined. A titer is calculated by multiplying the number of colonies produced for a given dilution by the dilution factor and is expressed as colony forming units (cfu) per μ l.

The library was plated in DNA format by inoculating 1000 cfu into 7 ml of semisolid LB-agar and grown for 48 hours at 30°C in 8-hole tubes (Autogen, Framingham, MA; cat no. AGP3210-10022) as described in the GeneTrapper™ protocol for semi-solid amplification of plasmid cDNA libraries (Life Technologies, Bethesda, MD). Colonies were harvested from the semisolid agar by centrifuging at 3000 rpm for 30 minutes. The agar was decanted and a conventional cosmid preparation procedure performed on an Autogen 740 robot (Autogen, Framingham, MA).

One hundred and twenty 8-hole tubes were inoculated in this way and ten 96-well DNA plates were generated. The ten DNA plates were ordered into a 2 x 5 matrix forming a matrix of 24 wells x 40 wells. 2 μ l from the well of every row of this 24 x 40 matrix were removed and pooled into 40 "row pools". Similarly 2 μ l from the well of every column were removed and pooled into 24 "column pools". Finally, 2 μ l from the well of every plate were removed and pooled into 10 "plate pools". In this way each well was represented in a "row pool", a "column pool", and a "plate pool". TE was added to each pool to bring the total volume to 150 μ l.

EXAMPLE 3 - Screening a pCosRKO Genomic Library

A) PCR-Screening Cosmid Pools

In order to screen for a GRI, at least a partial sequence for that gene should be known. From the available sequence information, PCR primers are designed to amplify a region of genomic DNA, preferably about 300 bp in size. A 1 μ l aliquot from each of the 74 row, column, and plate pools described above is used to PCR screen the pools in 96-well plate format using a conventional PCR amplification with

a Perkin Elmer GeneAmp® PCR System 9600 (Applied Biosystems Inc., Norwalk, CT). Positive row, column, and plate pools identified by this method are used to identify which of the 960 pools in the entire library of pools contain genomic clones of interest.

5 B) YTC amplification

Concurrent with library screening a YTC can be generated by PCR. A YTC is created by PCR amplification of a bacterial/mammalian positive selection marker using chimeric oligonucleotides that also bear identity to the gene of interest.

Bacterial/mammalian positive selection markers that are routinely used include the
 10 neomycin neo^r-cassette driven by dual bacterial/mammalian promoter in the plasmids pDual (Stratagene, La Jolla, CA) and a modified variant of pDual in which a NotI site has been inserted between the promoters and the neo^r coding sequence (pDual-NotI). The primers 44413-29FA (CCA TCA ACA CGC GTC TGC GTT CGA CCA GGC TGC GCG TTC TCG CGG CCA TAG CAA CCG ACG TAC GGC GTT GCG
 15 CCC TCG CCG GCA GCA AGA AGC CAC GGA AGT CAT GTG CGC GGA ACC CCT ATT TGT T; SEQ ID NO:10) and 44413-29RA(GAG CCA GAA CGG CGT CGG TCA CGG CAT AAG GCA TGC CCA TTG TTA TCT GGG CGC TTG TCA TTA CCA CCG CCG CGT CCC CGG CCG ATA TCT CAC CCT GGT CGA GAT AGC GCG GGT TCC TTC CGG TAT; SEQ ID NO:11) are used to amplify a neo^r-
 20 cassette from pDual-NotI. The 3' 23 base pairs of each of these primers bears identity to the 5' and 3' border of the pDual-NotI neo^r-cassette, respectively, while the 5' 103 base pairs of each oligonucleotide bear identity to regions in the hsvTK cassette of pCosRKO that we will call RA1 and RA2. Following PCR amplification the 1.6 kb product is gel isolated by running the PCR reaction out on a 0.8% TBE
 25 agarose gel and extracting the band using a Qiagen® gel extraction kit (Qiagen, Valencia, CA).

C) Yeast co-transfection of cosmid clone with YTC

For the purposes of an experimental control, an analogue of pCosRKO was made such that the regions RA1 and RA2 were deleted by cutting pCosRKO with
 30 the enzymes *AgeI* and *BspEI*, "polishing" the ends with T4 DNA polymerase, and religating the plasmid. This control plasmid was called pCosRKOdelta.

pCosRKODelta plasmid DNA was mixed with pCosRKO plasmid DNA in molar ratios of 1:1, 1:10, 1:100, and 1:1000. For each yeast transfection 100 ng of DNA from one of the above mixtures was used. To prepare a DNA mixture for transfection, DNA was linearized with the enzyme I-CeuI that cuts between two 50 base pair repeats in order to enhance transfection efficiency as described by Raymond et al. (BioTechniques, **27**:892-96 (1999)). The DGY63 yeast strain (R.D. Geitz, University of Manitoba) was transfected with 100 ng of I-CeuI digested pCosRKO and either 160 ng of the hsvTK-YTC described above according to the TRAF0 method described (Agatep et al., Technical Tips Online, (<http://tto.trends.com>) (1998)) with the exception that all solutions and media were made with ES Cell Qualified Ultra Pure H₂O (Specialty Media, Phillipsburg, NJ; cat # TMS-006-A). From the final suspension of transformed yeast in this protocol, 0.5 µl is diluted to 100 µl with sterile water and plated onto a 100 mm SC-Trp plate (Agatep et al., Technical Tips Online, (<http://tto.trends.com>) (1998)). In addition 5 µl, 50 µl, and 500 µl were diluted to 3 ml and plated onto 500 cm² SC-Trp plates (Agatep et al., Technical Tips Online, (<http://tto.trends.com>) (1998)). The 1.0 µl aliquots were used to deduce the yeast transformation efficiency per µg of DNA. From this efficiency the number of colonies on each of the 500 cm² plates was estimated. The yeast colonies were scraped off of each plate into 1-10 ml of YAPD media, spun down by centrifugation, and plasmid DNA prepared using ZymoprepTM (Zymo Research, Orange, CA). From the approximately 50 µl solution of DNA that was prepared, 3 µl was used to electroporate GeneHogs® electrocompetent bacterial cells (Research Genetics, Huntsville, AL). The electroporated bacteria were plated onto LB plates containing kanamycin (5 µg/ml). The number of kanamycin resistant clones was recorded. Table 1 shows the results of this experiment.

Table 1

Condition	Yeast cfu/plate	KAN cfu/plate	Transformation Efficiency
1:1	163	lawn	3.26 X10 ⁶
	1630	77	
	16,300	66	
	163,000	14	

1:10	438	119	8.76 X10 ⁶
	4380	50	
	43,800	0-arc	
	438,000	14-arc	
1:100	477	68	9.54 X10 ⁶
	4770	11	
	47,700	13	
	477,000	39	
1:1000	732	11	1.46 X 10 ⁶
	7320	0	
	73,200	39	
	732,000	296	

Since the Zymoprep™ method and bacterial electroporation are both known to be saturable it is not relevant to make predictions of homologous recombination frequency from the absolute *E. coli* kanamycin colony numbers. Instead we use a presence/absence call of kanamycin-resistant colonies from various yeast colony numbers to estimate homologous recombination frequency in yeast. Specifically, since kanamycin resistance was present 4 times out of 4 when the target plasmid was present amongst 100 fold excess of control plasmid, and 3 times out of 4 when the target plasmid was present amongst 1000-fold excess of control plasmid, we estimate that homologous recombination events were present at a frequency between 1 in 75 and 1 in 150, i.e., approximately 1 homologous recombination event in 112 yeast colonies. Fidelity of the recombination event has been verified by restriction analysis of plasmid DNA prepared from the kanamycin resistant colonies. Of 24 kan^r colonies isolated all demonstrated correct homologous recombination. We conclude that homologous recombination events reliably serve to identify rare plasmids in a mixture of other plasmids.

In the example shown above, the YTC was targeted into the TK cassette of the pCosRKO backbone. When preparing a targeting cassette for a GRI from a genomic library a YTC is directed to the GRI by making the appropriate chimeric oligonucleotides when PCR-amplifying the YTC. Since the library is divided into 960 pools of 1000 clones/pool, typically only one clone/pool or 1/1000th of the DNA

in a positive pool would be a target for the YTC. A Qiagen robot will yield approximately 3 µg of DNA for each pool. One tenth of this DNA (i.e., 300 ng) is used for the yeast electroporation along with 160 ng of the appropriate YTC for the GRI. This will yield at least 450,000 yeast transformants according to the yeast transformation efficiencies shown above. Of these, an estimated 450 will be transformed for the particular clone bearing the GRI. Based on the demonstrated homologous recombination frequency, approximately 4 will have undergone homologous recombination with the YTC and will be recovered as kanamycin resistant *E. coli* colonies. Plasmid DNA from any one of these colonies is suitable to be prepared and used for targeting ES cells.

EXAMPLE 4 - Characterizing Clones Isolated from a pCosRKO Genomic library

Individual clones identified as kanamycin resistant colonies on the above procedure are isolated and grown overnight in liquid culture at 30 °C in the presence of ampicillin (100 µg/ml) and plasmid DNA prepared by conventional mini-plasmid preparation methods known to those skilled in the art. This DNA is then restriction-mapped using the FLASH™ Nonradioactive Gene Mapping Kit (Stratagene Inc., La Jolla, Calif.) according to the procedures described by the manufacturer. Briefly this involves first completely digesting 10µg of the phage DNA with the restriction enzyme NotI using standard restriction enzyme digest conditions (Maniatis *et al.*, *supra*). NotI cuts all clones in the vector DNA at either end of the cloned insert, leaving a T3 bacteriophage promoter attached to one end of the insert and a T7 bacteriophage promoter attached to the other end. The NotI-digested DNA is then partially digested with the enzyme *EcoRI*, as an example, using limiting amounts of enzyme (0.2 units/µg DNA), in an 84 µl reaction volume at 37°C. Aliquots (26 µl) are removed after 3 minutes, 12 minutes, and 40 minutes, and the reaction is stopped by the addition of 1 µl of 0.5M EDTA. DNA from all three time points is resolved on a 0.7% agarose gel, visualized by ethidium bromide staining, and then transferred to a GeneScreen Plus® membrane (NEN Research Products, Boston, Mass.) by capillary transfer (Maniatis *et al.*, *supra*). The membrane is hybridized with an alkaline phosphatase labeled oligonucleotide that is specific for the T3 promoter (supplied with the FLASH™ kit) using reagents and methods supplied by the kit manufacturer. After hybridization, the membrane is washed and

developed with a chemiluminescent-yielding substrate and then exposed to X-ray film in the dark for approximately 60 minutes.

The oligonucleotide probes effectively label one end of the insert. By determining the positions of the bands on the X-ray film and calculating the DNA size to which they correspond, it is possible to determine the position of the *EcoRI* sites relative to the T3 end of the insert. These results are then complemented by stripping the probe off of the membrane, and rehybridizing with a T7-specific oligonucleotide in order to determine the positions of the *EcoRI* sites relative to the T7 end of the insert. This process is repeated using other 6 base pair-recognition site restriction endonucleases.

EXAMPLE 5 - Using a gene targeting vector generated from a pCosRKO genomic library

A) Embryonic Stem Cells

The DBA-252 ES cells are created as described (Roach, et al., Exp. Cell Res., 221:520-525, 1995). The cells are also grown as described therein.

B) DNA preparation

DNA is prepared for a gene targeting experiment by growing up a preparation of a gene targeting vector clone obtained by the above procedures and isolating DNA using a Qiagen® Maxi Kit (Qiagen, Valencia, CA). 400µg of targeting vector DNA are prepared for electroporation by linearizing the plasmid with either the enzyme *SwaI* or *PI-PspI* in a 1 ml reaction volume. The DNA is then precipitated by the addition of ethanol, washed with 70% ethanol, and resuspended in 500 µl of sterile water.

C) Electroporation of prepared targeting vector DNA into ES cells

The linearized targeting vector DNA is then electroporated into ES cells using a Bio-Rad Gene Pulser™ System (Bio-Rad Laboratories, Hercules, Calif.) as follows. In each of ten electroporation cuvettes, 40 µg of DNA is electroporated into 5×10^6 cells suspended in 0.8 ml ES cell medium. The electroporation conditions are 250 V and 500µF, which typically result in time constants ranging between 5.7-6.2 seconds. After electroporation the cells are incubated for 20 minutes at room temperature in the electroporation cuvettes. All the electroporated cells are then

pooled and distributed approximately equally onto 20 tissue culture plates (100 mm x 15 mm).

After 24 hours, the plates are aspirated and fresh ES cell medium is added. The following day, the medium in 19 plates is replaced with ES cell medium supplemented with 150 µg/mL of G418 (Life Technologies, Bethesda, MD) and 0.2 µM gancyclovir (Syntex, Palo Alto, CA). The medium in one plate is supplemented with 150µg/mL of G418 alone. After an additional 6 days, resultant individual ES cell colonies are picked off of the plates and separately expanded in individual wells of 24 well plates as described (Wurst and Joyner, Gene targeting, **126**:33-61, 1993). A comparison of the number of colonies that grow on the plates supplemented with G418 and gancyclovir versus the number that grow on the plates supplemented with G418 alone is used to determine the efficiency of negative selection, which typically between 3-10 fold.

D) Analyses of the gene-targeted ES cells

When the cell culture in each well of the 24-well plates becomes approximately 80% confluent, the cells are washed with PBS and then dispersed with two drops of trypsin-EDTA. Trypsinization is stopped by the addition of 1 ml of ES cell medium. An aliquot (0.5 mL) of this suspension is transferred to each of two wells of separate 24-well plates. After the cells have grown to near confluence, one of the plates is used for cryopreservation of the cell line while the other is used as a source of DNA for each of the cloned cell lines.

For cryopreservation, the cells in a 24-well plate are first chilled by placing the plate on ice. The medium is then replaced with fresh ES cell medium supplemented with 10% DMSO and 25% FBS. The plate is then cooled at approximately 0.5°C/minute by insulating the plate in a Styrofoam box and placing it in a -70 °C freezer.

To isolate the DNA from the cloned cell lines on the other 24-well plate, the medium in each well is replaced with 500 µl of digestion buffer (100 mM Tris-HCl, pH8.5, 5 mM EDTA, 0.2% SDS, 200 mM NaCl, and 100 µg/ml proteinase K) and incubated overnight at 37°C. After overnight incubation, 500µl of isopropanol is added to each well and the plate is agitated for 15 minutes on an orbital shaker. The supernatant fluid is aspirated and replaced with 500 µl of 70% ethanol and the

plate is shaken for an additional 15 minutes. The DNA precipitate is picked out of the well and dissolved in 50 µl of TE solution (10 mM Tris-HCl pH 7.5, 1 mM EDTA).

The analysis for a gene targeting event of the GRI involves a Southern hybridization screen of endonuclease digested ES cell DNA. The enzymes chosen for this procedure are ones that do not cut in one or the other homology arm of the gene targeting vector but do cut in the corresponding flanking MCS of the targeting arm. For example, an enzyme is chosen that does not cut in the left arm of the targeting vector but does cut in the MCS immediately flanking the left arm. Similarly, a second enzyme for a complementary digest would be chosen that does not cut in the right arm but does cut in the MCS immediately flanking the right arm. The probe for this analysis is derived from the bacterial/mammalian positive selection cassette. For the left arm analysis a probe is prepared by isolating the 350 base pair fragment obtained by digesting pDual-NotI with PvuII and NotI. The right arm probe is prepared by isolating the 660 base pair fragment obtained by digesting pDual-NotI with NotI and CspI.

An aliquot (10 µl) of each DNA sample is digested with the enzyme chosen for analysis of either the right or left arms, resolved on a 0.8% agarose gel, and transferred to a GeneScreen Plus™ membrane. The probe is prepared by first isolating the appropriate band from the double digested pDual-NotI (described above). The probe is labeled with ³²P-dCTP by random priming and hybridized overnight to the membrane at 58°C (Church *et al.*, Proc. Natl. Acad. Sci. USA, 81:1991-95, 1984). An ES cell line in which the GRI has been successfully targeted yields a fragment that is larger than that predicted by the restriction map of the gene targeting vector. This is the result of homologous recombination removing the MCS from the arm of homology as the targeting vector is incorporated into the genome by homologous recombination, whereas the MCS is not removed when the targeting vector randomly inserts into the genome.

All cell lines scored as putative homologous recombinants by this primary screen are then further screened using an aliquot of DNA digested with the appropriate enzyme to analyze the complementary arm of homology and probed with the alternative probe derived from pDual-NotI. Again, a bona fide homologous recombination event results in a band on the Southern blot that is larger than would be predicted from the restriction map of the gene targeting vector, due to the

removal of the MCS for that arm. Cells from the resulting cultures are used to make chimeric mice.

EXAMPLE 6 - Establishment of Gene-targeted mice homozygous for mutations in the GRI

Gene-targeted ES cells described above are used to make chimeric mice by aggregating the ES cells to E2.5 embryos and transferring the aggregated embryos to pseudopregnant females (See, e.g., Wood et al., *Nature*, **365**:87-89, 1993). ES cells are prepared for aggregation by limited trypsinization to produce clumps that averaged 10-15 cells. E2.5 embryos are collected from superovulated CD-1 female mice (albino) by oviduct flushing as described by Hogan et al. (*Manipulating the Mouse Embryo: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1986; incorporated herein by reference). The zona pellucida is removed from the embryos using acidic Tyrode's solution (Sigma Chemical Co., St. Louis, MO). Aggregation wells are created by pressing a blunt metal instrument (i.e., a darning needle) into tissue culture plastic. Embryos are then placed in a well together with a clump of approximately 10-15 ES cells in a small drop (approximately 20 μ l) of M16 medium (Sigma Chemical Co., St. Louis, MO) under mineral oil. After an overnight incubation (37 °C, 100% humidity, 5% CO₂ in air), approximately 20 of the aggregated embryos are transferred to the uterine horns of each pseudopregnant female (Hogan *et al.*, *supra*). Contribution of the ES cells to the offspring is scored by the appearance of pigmented coat color. Pigmented mice are termed chimeric founders. Germline contribution by the ES cells is scored by the appearance of pigmented offspring from a cross between the chimeric founders and CD-1 females.

The germline chimeras are then used to establish lines of mice bearing a copy of the mutation in the GRI. The presence of the gene-targeted mutation in the pigmented offspring is determined using the Southern blot strategy described above with genomic DNA prepared from a tail sample (Hogan *et al.*, *supra*). Mice that are homozygous for the mutation in the GRI are established by crossing 2 heterozygous mice and are identified using the same Southern blot strategy.

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention

5 has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the above-described modes for carrying out the invention which are obvious to those skilled in the field of molecular biology or related fields are intended to be within the scope of the

10 following claims.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2

Organization Applicant

Street : Eastern Point Road
City : Groton
5 State : CT
Country : USA
PostalCode : 06340-
PhoneNumber : 860-441-4901
FaxNumber : 860-441-5221
10 EmailAddress :
<110> OrganizationName : Pfizer Inc.

Individual Applicant

15 Street :
City :
State :
Country :
PostalCode : ____-____
20 PhoneNumber : ____-____-____
FaxNumber : ____-____-____
EmailAddress :
<110> LastName : Cole
<110> FirstName : Katherine
25 <110> MiddleInitial : E.
<110> Suffix :

Individual Applicant

30 Street :
City :
State :
Country :
PostalCode : ____-____
35 PhoneNumber : ____-____-____
FaxNumber : ____-____-____
EmailAddress :
<110> LastName : Reaume
<110> FirstName : Andrew
40 <110> MiddleInitial : G.
<110> Suffix :

Application Project

45 <120> Title : Rapid Creation of Gene Targeting Vectors Using
Homologous Recombination in Yeast
<130> AppFileReference : PC10667A
<140> CurrentAppNumber :
50 <141> CurrentFilingDate : ____-____-____

Sequence

55 <213> OrganismName : Primer
<400> PreSequenceString :
cgcttcgaaa tcgctgggcc attctcatga
30
<212> Type : DNA
<211> Length : 30
SequenceName : Sequence 1
60 SequenceDescription :

Custom Codon

Sequence Name : Sequence 1

5 Sequence

<213> OrganismName : Primer

<400> PreSequenceString :

10 cgcatttaaa ttcttccgct tcctcgctca ct
32

<212> Type : DNA

<211> Length : 32

SequenceName : Sequence 2

SequenceDescription :

15

Custom Codon

Sequence Name : Sequence 2

20 Sequence

<213> OrganismName : Primer

<400> PreSequenceString :

25 gaccgcttat catcgataag c
21

<212> Type : DNA

<211> Length : 21

SequenceName : Sequence 3

SequenceDescription :

30

Custom Codon

Sequence Name : Sequence 3

35 Sequence

<213> OrganismName : Primer

<400> PreSequenceString :

40 cgcatttaaa tgaaggctct caaggcatc gg
32

<212> Type : DNA

<211> Length : 32

SequenceName : Sequence 4

SequenceDescription :

45

Custom Codon

Sequence Name : Sequence 4

50 Sequence

<213> OrganismName : Primer

<400> PreSequenceString :

55 cgcggatcct agcctggcca tccgggccat cgctggggcca ttctcatga
49

<212> Type : DNA

<211> Length : 49

SequenceName : Sequence 5

SequenceDescription :

60

Custom Codon

Sequence Name : Sequence 5

Sequence

5

<213> OrganismName : Primer
<400> PreSequenceString :
cgcatcgatt tcgataagct ctgctttttg t
31

10

<212> Type : DNA
<211> Length : 31
SequenceName : Sequence 6
SequenceDescription :

15

Custom Codon

Sequence Name : Sequence 6

Sequence

20

<213> OrganismName : Primer
<400> PreSequenceString :
cgcttcgaac taccgggtag gggaggcgct
30

25

<212> Type : DNA
<211> Length : 30
SequenceName : Sequence 7
SequenceDescription :

30

Custom Codon

Sequence Name : Sequence 7

Sequence

35

<213> OrganismName : Primer
<400> PreSequenceString :
cgcagatcctt ccgatggccg cagcggcctc tgatggaatt agaacttggc a
51

40

<212> Type : DNA
<211> Length : 51
SequenceName : Sequence 8
SequenceDescription :

45

Custom Codon

Sequence Name : Sequence 8

50

Sequence

55

<213> OrganismName : Multiple Cloning Site
<400> PreSequenceString :
ggccgctgct tccaggctaa cacttatata ggcattggta ctgagataca gcgagatacc
60

gtgtaactat aacggtccta aggtagcgaa ggctaactat tatataggca ttggtactga
120

60

```

gatacagcga gataccgtgc ggccgctgcg gccaatatac cctcactaaa gggaggccta
180
tgcatactag tccatgggat atccccgggg ttaaccgcg gctcgagatc gatgaattcg
5 240
gccggccggt taaacggatc cggcgcgctt taattaaaag cttcatatgt ctagagtcga
300
cgagctcggg accgggcccc catgctggcc aaggcctaga tctccctata gtgagtcgta
10 360
ttatggcaaa cagctattat ggggtattat ggtaggccat ccggggccgcg gccgcttcca
15 420
tccggggc
428
<212> Type : DNA
<211> Length : 428
20 SequenceName : Sequence 9
SequenceDescription :

Custom Codon
-----
25 Sequence Name : Sequence 9

Sequence
-----
<213> OrganismName : Primer
30 <400> PreSequenceString :
ccatcaacac gcgtctgcgt tcgaccaggc tgcgcgttct cgcggccata gcaaccgacg
60
tacggcgttg cgccctcgcc ggcagcaaga agccacggaa gtcattgtgcg cggaaccctt
35 120
atttggtt
127
<212> Type : DNA
40 <211> Length : 127
SequenceName : Sequence 10
SequenceDescription :

Custom Codon
-----
45 Sequence Name : Sequence 10

Sequence
-----
50 <213> OrganismName : Primer
<400> PreSequenceString :
gagccagaac ggcgtcggtc acggcataag gcatgcccat tgttatctgg gcgcttgtca
60
ttaccaccgc cgcgtccccg gccgatatct caccctggtc gagatagcgc gggttccttc
55 120
cggtat
126
60 <212> Type : DNA
<211> Length : 126

```


References Cited

U.S. PATENT DOCUMENTS

5	5,777,195	7/1998.	Fienberg et al.....800/2
	5,814,300	9/1998.	Scott et al.424/9.1
	5,859,307	1/1999.	Mombaerts et al.800/2
	5,464,764	11/1995.	Capecchi et al.435/172.3
	5,789,215	9/ 1998.	Berns et al.....435/172.3
10	4,959,317	9/ 1990.	Sauer.....435/172.3

FOREIGN PATENT DOCUMENTS

WO 93/03183 11/1991 WIPO

OTHER PUBLICATIONS

- Agatep, R., R. D. Kirkpatrick, D. L. Parchaliuk, R. A. Woods, and R. D. Gietz.
Transformation of *Saccharomyces cerevisiae* by the lithium acetate/single-
stranded carrier DNA/polyethylene glycol (LiAc/ss-DNA/PEG) protocol.
Technical Tips Online (<http://tto.trends.com>), 1998.
- Audoly, L. P., S. L. Tilley, J. Goulet, M. Key, M. Nguyen, J. L. Stock, J. D. McNeish,
B. H. Koller, and T. M. Coffman. Identification of specific EP receptors
responsible for the hemodynamic effects of PGE2. *Am. J. Physiol.* 277: H924-
30, 1999.
- Capecchi, M. R. The new mouse genetics: Altering the genome by gene targeting.
Trends Genet 217: 70-76, 1989.
- Carlsson, L. M., J. Jonsson, T. Edlund, and S. L. Marklund. Mice lacking
extracellular superoxide dismutase are more sensitive to hyperoxia. *Proc. Natl.*
Acad. Sci 295: 6264-6268, 1995.
- Castle, W. E., and G. M. Allen. The heredity of albinism. *Proc. Am. Acad. Arts Sci.*
38: 603-621, 1903.

- Church, G. M., and W. Gilbert. Genomic sequencing. *Proc. Natl. Acad. Sci* 226: 1991-1995, 1984.
- Collins, F. S., A. Patrinos, E. Jordan, A. Chakravarti, R. Gesteland, and L. Walters. New goals for the U.S. Human Genome Project: 1998-2003. *Science* 282: 682-689, 1998.
- Doetschman, T., R. G. Gregg, N. Maeda, M. L. Hooper, D. W. Melton, S. Thompson, and O. Smithies. Targetted correction of a mutant HPRT gene in mouse embryonic stem cells. *Nature* 301: 576-578, 1987.
- Evans, M. J., and M. H. Kaufman. Establishment in culture of pluripotential cells from mouse embryos. *Nature* 309: 154-156, 1981.
- Gietz, R. D., and A. Sugino. New yeast - *Escherichia coli* shuttle vectors constructed with in vitro mutagenized yeast genes lacking six-base-pair restriction sites. *Gene* 74: 527-534, 1988.
- Hogan, B., F. Costantini, and E. Lacy. *Manipulating the mouse embryo: A Laboratory Manual*. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory, 1986.
- Koller, B. H., P. Marrack, J. W. Kappler, and O. Smithies. Normal development of mice deficient in beta 2M, MHC class I proteins, and CD8+ T cells. *Science* 248: 1227-1230, 1990.
- Koller, B. H., and O. Smithies. Altering genes in animals by gene targeting. *Ann. Rev. Immunol* 218: 705-30, 1992.
- Larionov, V., N. Kouprina, J. Graves, X. N. Chen, J. R. Korenberg, and M. A. Resnick. Specific cloning of human DNA as yeast artificial chromosomes by transformation-associated recombination. *Proc. Natl. Acad. Sci. USA* 93: 491-496, 1996.
- Li, Y., T.-T. Huang, E. J. Carlson, S. Melov, P. C. Ursell, J. L. Olson, L. J. Noble, M. P. Yoshimura, C. Berger, P. H. Chan, D. C. Wallace, and C. J. Epstein. Dilated cardiomyopathy and neonatal lethality in mutant mice lacking manganese superoxide dismutase. *Nature Genet.* 390: 376-381, 1995.
- Lowell, C. A., and P. Soriano. Genes Dev 1996 Aug 1;10(15):1845-57. *Genes Dev.* 10: 1845-1857, 1996.
- Maniatis, T., E. F. Fritsch, and J. Sambrook. *Molecular cloning: A laboratory manual*. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory, 1982.

- Martin, G. R. Isolation of a pluripotent cell line from early mouse embryos cultured in medium conditioned by teratocarcinoma stem cells. *Proc. Natl. Acad. Sci. USA* 78: 634-638, 1981.
- Mullis, K. B., and F. F.A. Specific synthesis of DNA in vitro via a polymerase-catalyzed chain reaction. *Methods Enzymol.* 155: 335-350, 1987.
- 5 Nagy, A., C. Moens, E. Ivanyi, J. Pawling, M. Gertsenstein, A. K. Hadjantonakis, M. Pirity, and J. Rossant. Dissecting the role of N-myc in development using a single targeting vector to generate a series of alleles. *Curr. Biol.* 8: 661-664, 1998.
- 10 Raymond, C. K., V. R. Mugford, and S. L. Sexson. Plasmid topologies that enhance the transformation efficiency of yeast. *BioTechniques* 27: 892-896, 1999.
- Reaume, A. G., P. A. deSousa, S. Kulkarni, B. L. Langille, D. Zhu, T. C. Davies, S. C. Juneja, G. M. Kidder, and J. Rossant. Cardiac malformation in neonatal mice lacking connexin43. *Science* 216: 1831-1833, 1995.
- 15 Reaume, A. G., J. L. Elliott, E. K. Hoffman, N. W. Kowall, R. J. Ferrante, D. F. Siwek, H. M. Wilcox, D. G. Flood, M. F. Beal, R. H. Brown. Jr., R. W. Scott, and W. D. Snider. Motor neurons in Cu/Zn superoxide dismutase-deficient mice develop normally but exhibit enhanced cell death after axonal injury. *Natur. Genet.* 13: 43-47, 1996.
- 20 Roach, M. L., J. L. Stock, R. Byrum, B. H. Koller, and J. D. McNeish. A new embryonic stem cell line from DBA/1lacJ mice allows genetic modification in a murine model of human inflammation. *Exp. Cell Res.* 221: 520-525, 1995.
- Skryabin, B. V., and C. Schmauss. Enhanced selection for homologous-recombinant embryonic stem cell clones with a neomycin phosphotransferase gene in antisense orientation. *Transgenic Res.* 6: 27-35, 1997.
- 25 Smithies, O., R. G. Gregg, S. S. Boggs, M. A. Koralewski, and R. S. Kucherlapat. Insertion of DNA sequences into the human chromosomal beta-globin locus by homologous recombination. *Nature* 317: 230-234, 1985..
- Spriggs, M. K., B. H. Koller, T. Sato, P. J. Morrissey, W. C. Fanslow, O. Smithies, R. F. Voice, M. B. Widmer, and M. C. R. Beta 2-microglobulin-, CD8+ T-cell-deficient mice survive inoculation with high doses of vaccinia virus and exhibit altered IgG responses. *Proc. Natl. Acad. Sci. USA* 89: 6070-6074, 1992.
- 30

- Sullivan, P. M., H. Mezdour, Y. Aratani, C. Knouff, J. Najib, R. L. Reddick, S. H. Quarfordt, and N. Maeda. Targeted replacement of the mouse apolipoprotein E gene with the common human APOE3 allele enhances diet-induced hypercholesterolemia and atherosclerosis. *J. Biol. Chem.* 272: 17972-17980, 1997.
- 5 Thomas, K., K. Folger and M. Capecchi. High frequency targeting of genes to specific sites in the mammalian genome. *Cell* 219: 419-428, 1986.
- Wood, S. A., W. S. Pascoe, C. Schmidt, R. Kemler, M. J. Evans, and N. D. Allen. Simple and efficient production of embryonic stem cell-embryo chimeras by coculture. *Proc. Natl. Acad. Sci.* 31: 4582-4585, 1993.
- 10 Wurst, W., and A. L. Joyner. Production of targeted embryonic stem cell clones. In: *Gene targeting*, edited by A. L. Joyner. Oxford, England: IRL Press at Oxford University Press, 1993, p. 33-61.
- Yang, D. D., C. Y. Kuan, A. J. Whitmarsh, M. Rincon, T. S. Zheng, R. J. Davis, P. Rakic, and R. A. Flavell. Absence of excitotoxicity-induced apoptosis in the hippocampus of mice lacking the Jnk3 gene. *Nature* 389: 865-870, 1997.
- 15

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50